

3.2 Synthetic Division

3.3 Zeros of Polynomial Functions

In these sections we will study polynomials algebraically. Most of our work will be concerned with finding the solutions of polynomial equations of any degree – that is, equations of the form

$$f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0 = 0 \quad (1)$$

Definition

A **root** or **solution** of equation (1) is a number k that when substituted for x leads to a true statement. Thus, k is a root of equation (1) provided $f(k) = 0$.

We also refer to the number k in this case as a **zero of the function f** .

Exercise #1

Checking for a zero or root.

a) Is -1 a zero of $P(x) = -x^3 + x^2 - x + 1$?

b) Is $x = \frac{1}{2}$ a root of the equation $2x^2 - 3x + 1 = 0$?

Note: If a root is repeated n times, we call it a **root of multiplicity n**

Exercise #2

a) State the multiplicity of each root of the equation: $x^2(x+1)^3(x-1) = 0$

b) Find all zeros and their multiplicities: $f(x) = 5x^2(x+1-\sqrt{2})(2x+5)$

c) Find all zeros and their multiplicities: $f(x) = (7x-2)^3(x^2+9)^2$

Division of Polynomials

The process of long division for polynomials follows the same four-step cycle used in ordinary long division of numbers: divide, multiply, subtract, bring down.

Notice that in setting up the division, we write both the dividend and divisor in decreasing powers of x .

Exercise #3 Divide $5x^3 - 6x^2 - 28x - 2$ by $x + 2$.

The result of the division can be written as: _____

or

Note 1) Second equation is valid for all real numbers x , whereas first equation carries implicit restrictions that x may not equal -2 . For this reason, we often prefer to write our results in the form of the second equation.

2) The degree of the remainder is less than the degree of the divisor. This is very similar to the situation with ordinary division of positive integers, where the remainder is always less than the divisor.

The Division Algorithm

Let $f(x)$ and $g(x)$ be polynomials with $g(x)$ of lower degree than $f(x)$ and assume that $g(x) \neq 0$. Then there are unique polynomials $q(x)$ and $r(x)$ such that

$$f(x) = g(x) \cdot q(x) + r(x)$$

where $r(x) = 0$ or the degree of $r(x)$ is less than the degree of $g(x)$.

The polynomials $f(x)$ and $g(x)$ are called the **dividend** and **divisor**, respectively, $q(x)$ is the **quotient**, and $r(x)$ is the **remainder**.

When $r(x) = 0$, we have $f(x) = g(x) \cdot q(x)$ and we say that $g(x)$ and $q(x)$ are **factors** of $f(x)$.

Exercise #4 Using long division to find a quotient and a remainder.

Divide $x^3 + 2x^2 - 4$ by $x - 3$.

Synthetic Division

- Synthetic division is a quick method of dividing polynomials.
- It can be used **when the divisor is of the form $x - k$** .
- In the synthetic division we write down only the essential parts of the long division table (the coefficients).

Exercise #5 Use synthetic division to perform the following divisions:

a)
$$\frac{x^3 + 4x^2 - 5x + 44}{x + 6}$$

If $f(x) = x^3 + 4x^2 - 5x + 44$, evaluate $f(-6)$. What do you observe?

$$\text{b) } \frac{\frac{1}{3}x^3 - \frac{2}{9}x^2 + \frac{1}{27}x + 1}{x - \frac{1}{3}}$$

If $f(x) = \frac{1}{3}x^3 - \frac{2}{9}x^2 + \frac{1}{27}x + 1$, evaluate $f\left(\frac{1}{3}\right)$. What do you observe?

The Remainder Theorem

When we divide a polynomial $f(x)$ by $x - k$, the remainder is $f(k)$.

Proof

Exercise #6 Using the remainder theorem to evaluate a function and check for a factor.

a) Let $f(x) = x^2 + 5x + 6$.

i) Evaluate $f(-2)$.

ii) Is $x + 2$ a factor of $f(x) = x^2 + 5x + 6$?

b) Let $f(x) = 6x^4 + x^3 - 8x^2 + 5x + 6$.

i) Evaluate $f\left(\frac{1}{2}\right)$.

ii) Is $x - \frac{1}{2}$ a factor of $f(x) = 6x^4 + x^3 - 8x^2 + 5x + 6$?

The Factor Theorem

The polynomial $x - k$ is a factor of the polynomial $f(x)$ if and only if $f(k) = 0$.

Exercise #7

Let $f(x) = 2x^3 - 4x^2 + 2x - 1$.

a) What is the remainder when dividing the given polynomial by $x - 2$? In how many ways can you find the remainder? Which method is the easiest one?

b) Is $x - 2$ a factor of $f(x)$?

c) Is $x - 1$ a factor of $f(x)$?

Exercise #8 Factoring a polynomial given a zero.

a) Let $f(x) = 6x^3 + 13x^2 - 14x + 3$. Show that -3 is a zero and use this fact to factor $f(x)$ completely.

b) $f(x) = 2x^4 + x^3 - 9x^2 - 13x - 5$. Knowing that -1 is a root of multiplicity 3, factor $f(x)$ into linear factors.

The Conjugate Zeros Theorem

If $f(x)$ is a polynomial function with real coefficients and if $a + bi$ is a zero of $f(x)$, then its conjugate $a - bi$ is also a zero of $f(x)$.

Exercise #9 For each polynomial, one zero is given. Find all the others.

a) $f(x) = x^3 - 7x^2 + 17x - 15$; $2 - i$

b) $f(x) = 4x^3 + 6x^2 - 2x - 1; \frac{1}{2}$

The Fundamental Theorem of Algebra

Every polynomial equation of the form

$$f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0 = 0 \quad (n \geq 1, a_n \neq 0)$$

has at least one root within the complex number system. (This root may be a real number).

The Linear Factors Theorem

Every polynomial of degree n can be expressed as a product of n linear factors.

$$f(x) = a_n (x - x_1)(x - x_2) \dots (x - x_n),$$

where a_n is the leading coefficient and x_i are the roots of the polynomial.

Theorem

Every polynomial of degree $n \geq 1$ has exactly n roots, where a root of multiplicity k is counted k times.

Exercise #10 Write each polynomial as a product of linear factors.

a) $f(x) = 3x^2 - 5x - 2$

b) $f(x) = x^2 - 5$

c) $f(x) = x^2 - 4x + 5$

Exercise #11 Finding polynomial equations satisfying given conditions.

In each case, find a polynomial equation $f(x) = 0$ satisfying the given conditions. If there is no such equation, say so.

a) Find a polynomial function of degree 3 having the numbers -3, 1, and 4 as roots and satisfying $f(2) = 30$.

b) A factor of $f(x)$ is $x-3$, and -4 is a root of multiplicity 2.

c) Find a polynomial function of degree 3 having the number -3 as a zero of multiplicity 3 and satisfying the condition $f(3) = 36$.

Exercise #12 Find a polynomial function of least degree having only real coefficients with zeros as given. What is the degree of each polynomial?

a) 2 and $1+i$.

b) $5 + i$ and $4 - i$.

The Number and Location of Real Zeros

Descartes' Rule of Signs

In some cases, the following rule – discovered by the French philosopher and mathematician Rene Descartes around 1637 – is helpful in eliminating candidates from lengthy lists of possible rational roots.

To describe this rule, we need the concept of **variation in sign**. If $f(x)$ is a polynomial with real coefficient, written with descending powers of x (and omitting powers with coefficient 0), then a variation in sign is a change from positive to negative or negative to positive in successive terms of the polynomial (adjacent coefficients have opposite signs).

Example #5 How many variations in sign occur in the following polynomial?

$$f(x) = 5x^7 - 3x^5 - x^4 + 2x^2 + x - 3$$

Descartes' Rule of Signs

Let $f(x)$ be a polynomial with real coefficients and a nonzero constant term.

- a) The number of positive real zeros of $f(x)$ is either equal to the number of variations in sign in $f(x)$ or is less than that by an even whole number.
- b) The number of negative real zeros of $f(x)$ is either equal to the number of variations in sign in $f(-x)$ or is less than that by an even whole number.

Exercise #15 For each polynomial function

- i) List the maximum number of real zeros;
- ii) List the number of positive real zeros and negative real zeros;
- iii) list all possible rational zeros;
- iv) find all rational zeros;
- v) factor $f(x)$.

a) $f(x) = x^3 + 6x^2 - x - 30$.

b) $f(x) = 15x^3 + 61x^2 + 2x - 8$

Exercise #16 a) Find all the complex zeros of $f(x) = x^4 - 6x^3 + 22x^2 - 30x + 13$.

b) Find all the solutions of $x^4 - 5x^3 - 5x^2 + 23x + 10 = 0$.